

# Oil sands development contributes polycyclic aromatic compounds to the Athabasca River and its tributaries

Erin N. Kelly<sup>a</sup>, Jeffrey W. Short<sup>b</sup>, David W. Schindler<sup>a,1</sup>, Peter V. Hodson<sup>c</sup>, Mingsheng Ma<sup>a</sup>, Alvin K. Kwan<sup>a</sup>, and Barbra L. Fortin<sup>a</sup>

<sup>a</sup>Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada T6G 2E9; <sup>b</sup>Oceana, 175 South Franklin Street, Suite 422, Juneau, Alaska 99801; and <sup>c</sup>Department of Biology and School of Environmental Studies, Queen's University, Kingston, ON, Canada K7L 3N6

Contributed by David W. Schindler, October 23, 2009 (sent for review September 21, 2009)

For over a decade, the contribution of oil sands mining and processing to the pollution of the Athabasca River has been controversial. We show that the oil sands development is a greater source of contamination than previously realized. In 2008, within 50 km of oil sands upgrading facilities, the loading to the snowpack of airborne particulates was 11,400 T over 4 months and included 391 kg of polycyclic aromatic compounds (PAC), equivalent to 600 T of bitumen, while 168 kg of dissolved PAC was also deposited. Dissolved PAC concentrations in tributaries to the Athabasca increased from 0.009  $\mu\text{g/L}$  upstream of oil sands development to 0.023  $\mu\text{g/L}$  in winter and to 0.202  $\mu\text{g/L}$  in summer downstream. In the Athabasca, dissolved PAC concentrations were mostly <0.025  $\mu\text{g/L}$  in winter and 0.030  $\mu\text{g/L}$  in summer, except near oil sands upgrading facilities and tailings ponds in winter (0.031–0.083  $\mu\text{g/L}$ ) and downstream of new development in summer (0.063–0.135  $\mu\text{g/L}$ ). In the Athabasca and its tributaries, development within the past 2 years was related to elevated dissolved PAC concentrations that were likely toxic to fish embryos. In melted snow, dissolved PAC concentrations were up to 4.8  $\mu\text{g/L}$ , thus, spring snowmelt and washout during rain events are important unknowns. These results indicate that major changes are needed to the way that environmental impacts of oil sands development are monitored and managed.

airborne deposition | oil sands processing | water contamination | hydrocarbons | oil sands mining

The Alberta oil sands consist of water, sand, and bitumen, a heavy and viscous hydrocarbon, that is recovered by surface mining or by in situ steam injection. To produce crude oil, bitumen must be extracted with hot water and upgraded by using heat, pressure, and catalysts (1). Production of bitumen increased from 482,000 to 1.3 million barrels per day from 1995 to 2008 (2, 3). The area disturbed by mine operations was 530 km<sup>2</sup> in 2007, and the area of tailings ponds surpassed 130 km<sup>2</sup> in 2008 (1, 4). Oil sands production by both mining and in situ methods will increase rapidly, with projected output ranging from 2.0 to 2.9 million barrels per day by 2020 (5).

Some residents of downstream Fort Chipewyan are convinced that the oil sands industry is responsible for higher than expected cancer rates (6). However, government, industry and related agencies, relying in part on the joint Regional Aquatic Monitoring Program (RAMP), report that effects are minimal, that natural sources cause elevated contaminant concentrations in the Athabasca and its tributaries (7), and that human health and the environment are not at risk from oil sands development (8–10), see *Controversy Background Information* in *SI Text*.

Since 1997, the RAMP, funded by industry and directed by a multistakeholder committee, has monitored aquatic ecosystems near the oil sands development (11). However, it lacks scientific oversight, and a peer review severely criticized its ability to detect effects (12). RAMP data are not publicly available, and the methods used to analyze, interpret, and report the data are not entirely transparent.

We conducted an independent, detailed, and accessible assessment of the loadings of polycyclic aromatic compounds (PAC) to the north-flowing Athabasca River, its tributaries, the Athabasca Delta, and Lake Athabasca (Fig. 1). In February to March and June to August 2008, we sampled water using polyethylene membrane devices (PMDs). In March, the accumulated snowpack was sampled at most sites. Athabasca River sites were selected upstream and downstream of oil sands mining and processing activity. Upstream and downstream sites near oil sands development are directly exposed to erosion of the McMurray geologic formation (McMF), where most oil sands occur (13).

Three sites along each of four impacted tributaries were selected using 2006 Landsat imagery. The first was located upstream of oil sands development and the McMF, the second midstream within the McMF but upstream of mining activity, and the third near stream mouths at the confluence with the Athabasca, downstream of development and downstream or within the McMF. Comparable sites were chosen on two reference tributaries unaffected by industry. In the summer, additional stream mouth sites, with and without upstream development, were included to increase statistical power.

After sampling, 2008 Landsat imagery revealed marked changes in the extent of oil sands development since 2006. Some tributary sites could be compared as planned, but some midstream and stream mouth sites unaffected in 2006 were affected by new development in 2008. To assess the impacts of this new development, the change in development between 2006 and 2008 was categorized visually from Landsat imagery as nil–small (N-S) and medium–large (M-L). This gave four classes of new development: midstream/N-S, midstream/M-L, stream mouth/N-S, and stream mouth/M-L. To compare the relative importance of natural erosion and mining on PAC mobilization, PAC concentrations in water were regressed against the proportion of the catchment within the McMF, overall land disturbance, and land disturbed by oil sands mining in 2008. These comparisons were made for all tributaries combined and separately for the Athabasca.

Samples were analyzed for PAC (sum of parent and alkylated homologues of two-, three- and four- ring polycyclic aromatic hydrocarbons + dibenzothiophene). Melted snow was analyzed for the mass of particulate and associated PAC retained on 0.45- $\mu\text{m}$  glass fiber filters and for dissolved PAC in filtrate. Because PMDs accumulate only dissolved PAC from water (14), equivalent water concentrations were calculated from PAC concentrations in PMDs,


Author contributions: E.N.K., J.W.S., D.W.S., and P.V.H. designed research; E.N.K., M.M., A.K.K., and B.L.F. performed research; E.N.K. and J.W.S. analyzed data; and E.N.K., J.W.S., D.W.S., and P.V.H. wrote the paper.

The authors declare no conflict of interest.

Freely available online through the PNAS open access option.

<sup>1</sup>To whom correspondence should be addressed. E-mail: d.schindler@ualberta.ca.

This article contains supporting information online at [www.pnas.org/cgi/content/full/0912050106/DCSupplemental](http://www.pnas.org/cgi/content/full/0912050106/DCSupplemental).



**Fig. 1.** Study Area Map. Sites: blue, Athabasca River; black, tributaries (AR17U, unnamed creek; AR17D, unnamed creek; HOR, Horse River; CLCR, Clarke Creek; POP, Poplar Creek, BE, Beaver River; ST, Steepbank River; MCC, McLean Creek; MACK, MacKay River; EL, Ells River; JOC, Jocelyn Creek; MU, Muskeg River; FR, Firebag River; FOR, Fort Creek; TR, Tar River; CALR, Calumet River; EYC, Eymundson Creek; 1, upstream; 2, midstream; 3, stream mouth). Landsat 5 image is a false color composite, where blue is water, green is vegetation, and pink is nonvegetated and/or developed areas. Squares represent existing and approved oil sands projects.

assuming equilibrium between PMDs and ambient water concentrations after  $\approx 30$ -day deployment. Further details are provided in *Methods* and *Analytical and Statistical Method Details* in *SI Text*.


## Results

**Particulates and PAC in Snow.** Substantial deposition of airborne particulates was discovered within 50 km of the Suncor and Syncrude upgrading facilities, near AR6 (Fig. 2 and Fig. S1). Particulate deposition exponentially declined from  $19 \text{ g/m}^2$  at AR6, near the upgrading facilities, to  $<0.35 \text{ g/m}^2$  at sites  $>50 \text{ km}$  distant (particulates =  $10.6 \times e^{-0.0714 \cdot x} \text{ g/m}^2$ ,  $x = \text{km}$  from AR6,  $r^2 = 0.71$ ,  $P < 0.0001$ ,  $n = 23$ ; Fig. S2A). Integrating over a 50-km radius indicates deposition of 11,400 metric T of particulates during  $\approx 4$  months of snowfall.


Most particulates collected at AR6 consisted of oil sands bitumen. An oil slick formed on the surface of melted snow (Fig. S3), and the PAC distribution was similar to the four oil sands samples, i.e., dominated by dibenzothiophenes, phenanthrenes/anthracenes, fluoranthenes/pyrenes, and benzantracenes/chrysenes (Fig. S4 A and B). Compared with the oil sands, PAC in the snow particulates were slightly enriched in naphthalenes and unsubstituted four- to five-ring PAC, suggesting admixture with volatile and combustion-derived PAC [i.e., dominated by less-substituted four or more-ring PAC (15)]. Using a measured mean ratio of  $0.000649 \pm 0.000168$  (see *Analytical and Statistical Method Details* in *SI Text*) for total PAC to bitumen in oil sands, the 0.414 mg of PAC per gram of particulate in AR6 snow was equivalent to 64% bitumen.

The proportion of PAC in snow particulates declined rapidly with distance from AR6, indicating that bitumen was deposited closer to the source than particulates (Fig. 3). Measured PAC deposition declined exponentially (PAC =  $1.06 \times e^{-0.130 \cdot x} \text{ mg/m}^2$ ,  $r^2 = 0.76$ ,  $P < 0.0001$ ,  $n = 23$ ; Fig. S2B) from  $7.87 \text{ mg/m}^2$  at AR6, near the upgrading facilities, to  $0.011 \text{ mg/m}^2$  or less at sites over 50-km distant. This is equivalent to deposition of 391 kg of PAC within a 50-km radius of AR6 over 4 months, or 600 T of bitumen. Deposition of PAC declined significantly more rapidly with distance from AR6 than particulates ( $t$  test,  $P < 0.01$ ), suggesting association of bitumen PAC with heavier or denser particles.

Dissolved PAC in the filtrate from melted snow also declined with distance from AR6 (Fig. 3), and were dominated by fluorenes, dibenzothiophenes, and phenanthrenes/anthracenes, consistent with partitioning into air or water of more volatile and soluble PAC in bitumen (Fig. S4C). Dissolved PAC deposition declined exponentially with distance from AR6 (dissolved PAC =  $0.148 \times e^{-0.0691 \cdot x} \text{ mg/m}^2$ ,  $r^2 = 0.59$ ,  $P < 0.0001$ ,  $n = 23$ ; Fig. S2C), with a similar decay constant ( $-0.0691$ ) as particulates ( $-0.0714$ ;  $P > 0.8$ ). Integrating over a 50-km radius gives 168 kg of dissolved PAC accumulated during 4 months. Dissolved PAC concentrations in



**Fig. 2.** White  $0.45\text{-}\mu\text{m}$  Whatman GF/F filters after 900 mL of melted snow from each site was filtered. Yellow, distance between sites; blue, communities; red, existing and approved surface mining projects.




**Fig. 3.** Particulate and dissolved PAC loading ( $\text{mg}/\text{m}^2$ ) in accumulated snowpack, March 2008. (A) Athabasca River, Athabasca Delta, and Lake Athabasca. (B) Tar River. (C) Joslyn Creek. (D) Ells River. (E) Beaver River. (F) Firebag River. (G) Muskeg River. (H) Steepbank River. Bolded font indicates existing and approved projects (Syncrude: Mildred Lake and Aurora, Suncor: Millennium, Steepbank and Firebag In-situ, Petro Canada Dover-MacKay River In-situ, Husky Oil Sunrise In-situ, TOTAL/Deer Creek In-situ, Shell: Albian Sands/Muskeg River and Jackpine, CNRL: Horizon, Imperial Oil: Kearl, Petro Canada/UTS: Fort Hills). Mildred Lake spans AR16 to AR7, whereas Steepbank and Millennium span AR16 to AR6, as indicated by arrows. Upgrading facilities are near AR6.

melted snow declined from  $4.8 \mu\text{g}/\text{L}$  at AR6 to  $<0.27 \mu\text{g}/\text{L}$  at sites  $>50 \text{ km}$  away, and exceeded  $0.7 \mu\text{g}/\text{L}$  at 9 of 10 sites within 22 km of AR6.

**PAC in Tributaries.** Dissolved PAC concentrations in the six tributaries sampled during winter and summer mostly increased

from upstream to downstream and were greater in summer than winter (Fig. 4A). Mean concentrations increased from  $0.009 \mu\text{g}/\text{L}$  at upstream sites in both winter and summer to  $0.023 \pm 0.0059 \mu\text{g}/\text{L}$  and  $0.202 \pm 0.160 \mu\text{g}/\text{L}$  at stream mouth sites, respectively, similar to melted snow. Differences among sites along tributaries were highly significant (two-way ANOVA, In



**Fig. 4.** Estimated winter and summer dissolved PAC concentrations ( $\mu\text{g}/\text{L}$ ). (A) Tributaries. (B) Midstream and stream mouth tributary sites in relation to “new” development (change in development from 2006 to 2008,  $n = 5-8$ . Athabasca River, Athabasca Delta, and Lake Athabasca (C) Winter. (D) Summer. \* indicates samples contaminated by diesel fuel (see *Analytical and Statistical Method Details* in *SI Text*), not included in calculations; McMf, McMurray Formation; N, north. Error bars are standard error of the mean.

transformed,  $P = 0.004$ ), but not between seasons ( $P = 0.91$ ). The power to detect seasonal differences was low ( $\beta = 0.05$ ), so a seasonal effect could not be discounted. The greater trend of increasing PAC downstream in summer compared with winter (Fig. 4A) was nearly significant (interaction  $P = 0.095$ ) at only moderate power ( $\beta = 0.30$ ).

This analysis included all upstream, midstream, and downstream tributary sites, regardless of development. Almost all development was near the stream mouth on some tributaries, with insignificant development near midstream or upstream sites. Analyzing only these tributaries, no significant increase in PAC concentrations between upstream and midstream sites was found (Fig. S5A). Thus, when development was insignificant, flow of water through the McMF did not significantly affect PAC concentrations, indicating that natural sources are not solely responsible for increased concentrations of PAC midstream or at stream mouths.

Increasing PAC concentrations from upstream to downstream in the tributaries (Fig. 4A and Fig. S5A) could reflect increasing contributions from natural erosion of the McMF, greater disturbance from development, or both. Little of the variability in summer PAC concentrations of tributaries was explained by the proportion of McMF, total surface land disturbance, or oil sands mining disturbance in watersheds ( $r^2_{\text{McMF}} = 0.009$ ,  $r^2_{\text{disturbance}} = 0.003$ ,  $r^2_{\text{oil sands}} = 0.000$ ;  $P > 0.69$ ,  $df = 19$ ).

In contrast, when midstream and stream mouth sites were grouped, PAC concentrations were strongly associated with new land disturbance or mining activity expansion from 2006 to 2008. Mean PAC concentrations increased 2-fold from  $0.012 \pm 0.0012$  at N-S sites to  $0.027 \pm 0.0052$   $\mu\text{g/L}$  at M-L sites in winter, and 8-fold from  $0.024 \pm 0.0055$  to  $0.197 \pm 0.0738$   $\mu\text{g/L}$  in summer (Fig. 4B). Season and disturbance effects were highly significant (two-way ANOVA, ln transformed,  $P < 0.004$ ), and the season and disturbance interaction was nearly significant ( $P = 0.065$ ,  $\beta = 0.36$ ). The power to detect the seasonal effect was  $\beta > 0.85$ , much greater than for the previous test ( $\beta = 0.05$ ). Comparing only midstream or stream-mouth sites leads to essentially the same conclusions (Fig. S5B and C). Thus, seasonal differences in PAC concentrations are likely real, and increased PAC concentrations in both winter and summer result from land disturbance by oil sands development between 2006 and 2008.

Approximately 75% of PAC homologues accumulated by PMDs deployed at the M-L sites consisted of three-ring PAC, dominated by alkyl-substituted dibenzothiophenes, phenanthrenes/anthracenes and fluorenes, with the remainder mostly four-ring PAC including alkyl-substituted fluoranthenes/pyrenes and benzanthracenes/chrysenes and negligible naphthalenes (Fig. S4D). Enrichment of three-ring PAC in PMDs compared with oil sands (Fig. S4A) is consistent with greater solubility of three- vs. four-ring PAC. The near absence of naphthalenes in PMDs is also consistent with low concentrations of these PAC in bitumen (Fig. S4A).

**Athabasca River, Athabasca Delta, and Lake Athabasca.** Dissolved PAC concentrations were usually low at most sites on the Athabasca River, Athabasca Delta, and Lake Athabasca, but often greater during summer (Fig. 4C and D). In winter, concentrations were mostly  $< 0.025$   $\mu\text{g/L}$ , except at sites near oil sands upgrading facilities and tailings ponds which ranged from 0.031 to 0.083  $\mu\text{g/L}$  (Fig. 4C). In summer, PAC concentrations in the Athabasca were usually  $< 0.030$   $\mu\text{g/L}$ . Upstream and within oil sands development, concentrations were unrelated to the proportion of McMF, total surface land disturbance, and oil sands mining disturbance ( $r^2_{\text{McMF}} = 0.137$ ,  $r^2_{\text{disturbance}} = 0.006$ ,  $r^2_{\text{oil sands}} = 0.085$ ;  $P > 0.33$ ,  $df = 8$ ). However, immediately downstream of new development, concentrations ranged from 0.063 to 0.135  $\mu\text{g/L}$  (Fig. S6).

## Discussion

The increased deposition of particulates and PAC in snow close to the Suncor and Syncrude upgrading facilities clearly implicates

them as sources and corresponds to a similarity between patterns of PAC congeners in particulates and oil sands (Fig. S4A and B). The enrichment of snow particulates by the more volatile PAC and by five-ring PAC (e.g., benzo[a]pyrene) is typical of PAC volatilized by heat or particulates produced by combustion. The dominance of oily material in snow from AR6 also suggests a separate organic bitumen phase in stack emissions that is present as droplets larger and less buoyant than average particulates, and precipitates near the source. Alternatively, the heavier particles might be bitumen-contaminated dust eroded by wind from mine sites (Fig. S7), but this was inconsistent with the high organic content of particulates near AR6. Although mining can mobilize dust, deposition would likely be localized and site-specific, and further study is needed to establish detailed loadings.

The similar deposition patterns of particulates and dissolved PAC in melted snow (Fig. S2A and C) suggests that dissolved PAC did not leach from particulates, but was scavenged from the atmosphere. Snow samples were filtered within an hour of thawing, leaving little time for PAC dissolution. More likely, PAC dissolved in the snowmelt were readily desorbed from non-bitumen particulates or scavenged from vapor-phase PAC by ice nuclei in plumes of condensing steam from stack emissions, as occurs with metals (16).

The oil sands industry is a known source of air pollutants. Snow surveys in 1978 and 1981 identified elevated metal deposition via flyash particulates 25 km north and south, and 10 km east and west of Suncor and Syncrude upgrading facilities (16, 17). In 1978, 96% of particulates were deposited within 25 km of the stacks (17), but PAC deposition was not measured. From 2005 to 2007, the mean annual release of particulates measured by Suncor and estimated by Syncrude (stack and fugitive emissions) was  $6037 \pm 927$  T (18). In contrast, during the 4 months before sampling in 2008, emissions were almost twice as large at 11,400 T and contaminated an area nearly 2-fold larger, with only  $\approx 60\%$  of particulates falling within a 25-km radius. Assuming similar deposition rates during the year implies a total annual particulate deposition of  $\approx 34,000$  T. This is nearly five times current reported emissions, and similar to annual deposition rates of 32,594 T in 1978 (19), before precipitators were installed. The discrepancy may be due to dust from mining (Fig. S7) or somewhat elevated loading estimates that were based on a circle around AR6, despite somewhat greater particulate deposition north/south of upgrading facilities than east/west (16). The close association of deposition with proximity to the upgrading facilities suggests that they are the primary source.

Airborne PAC from oil sands development conveys a considerable burden to the surrounding watershed. Historical stack discharges of particulates rich in aluminum (Al) (16, 17), and a strong correlation between Al and PAC concentrations in snow ( $r = 0.94$ ,  $P < 0.001$ ), suggest that large amounts of particulate PAC have been discharged since the onset of oil sands production in the 1960s. If deposition rates are constant throughout the year, the estimated annual release of PAC is now  $\approx 1,200$  kg associated with  $\approx 1,800$  T of bitumen particulates, and another 500 kg of dissolved PAC. This amount of bitumen released in a pulse would be equivalent to a major oil spill, repeated annually.

Given that particulate deposition rates in the 1970s before installation of stack precipitators (19) were as great as today, this situation has likely persisted for 30–40 years. As a result, current background PAC concentrations in surface soils, vegetation, snow, and runoff over a broad area of boreal forest may be greater than true background concentrations contributed naturally by oil sands in the region. Although RAMP collects snow for hydrologic monitoring (11), pollutant concentrations are not reported. In the early 1980s, snow sampling was recommended in northeastern Alberta and adjacent areas of Saskatchewan and the Northwest Territories to assess the effects of air emissions from expanding oil sands development (16, 17). The absence of such a program has made it progressively more difficult to separate pollution inputs

from rising background contamination. With more oil sands development projects approved and proposed, including new and expanded upgrading facilities, the increased deposition of airborne PAC will further raise regional “background” concentrations.

Tributaries impacted by oil sands development indicate a second major flux of PAC to receiving waters. Recent disturbances (new roads, deforestation, encampments, exploration, mining) expose and distribute fresh bitumen to wind and soil erosion and enhance bitumen transport to surface waters, which leach out the most available PAC. The lack of correlation between PAC concentrations in water and the extent of older development suggests that disturbed areas eventually stabilize. However, the lack of correlation may also reflect increasing background PAC concentrations. At sites distant from upgrading facilities and unaffected by land disturbance, the regional background of total dissolved PAC in surface waters is  $\approx 0.015 \mu\text{g/L}$ , closely comparable with concentrations in remote Canadian Arctic rivers (20). In contrast, at the most impacted stream mouths, PAC concentrations were 10- to nearly 50-fold greater (e.g., EL3, PAC =  $0.682 \mu\text{g/L}$ ), similar to concentrations toxic to fish embryos [as low as  $0.4 \mu\text{g/L}$  (21)]. The PAC in oil sands, snow, and water were dominated by homologues of three-ringed alkyl phenanthrenes, alkyl dibenzothiophenes, and alkyl fluorenes, PAC most closely associated with embryotoxicity of crude oil (22, 23). Embryos of fathead minnows (*Pimephales promelas*) and white sucker (*Catostomus commersoni*), species native to the Athabasca watershed, showed higher rates of mortality, reduced rates of growth, and signs of pathology typical of PAC toxicity when exposed to as little as  $0.01\text{--}0.1 \mu\text{g/L}$  of alkyl phenanthrene in oil sands leachates (calculated from refs. 24 and 25). PAC can also limit fish production through endocrine disruption. Compared with reference fish, gonads of slimy sculpin (*Cottus cognatus*) and pearl dace (*Margariscus margarita*) collected near active oil sands processing were less capable of synthesizing sex steroids (26). PAC may contribute to a greater prevalence of abnormal juvenile and adult fish captured in the Athabasca near and downstream of oil sands mining (11, 27).

During spring, the snowmelt pulse could increase PAC concentrations in tributaries to those toxic to both aquatic and terrestrial organisms (28, 29). Dissolved PAC would be immediately available to biota but particulate PAC may be taken up by filter feeders or partition into water for uptake by fish across gills. Residual particles may also accumulate on the forest floor with organic material or in underlying soils, causing PAC to leach more gradually into surface waters, adding to the overall PAC burden.

Of the 24 fish species resident in the Athabasca and its tributaries (30), 19 spawn in the spring or early summer (31). Embryos of these species are likely present when PAC concentrations are greatest. If located in shallow tributaries receiving PAC-enriched snowmelt, embryos may also experience photo-enhanced toxicity (32). Newly hatched whitefish embryos exposed to sublethal concentrations of retene (alkyl phenanthrene) died when coexposed to visible and UV light (33). Toxicity may also increase if PAC and metals associated with oil sands act synergistically, as observed for *Daphnia magna* (34).

Dissolved PAC did not persist as far as the Athabasca River Delta and Ft. Chipewyan, at least during the seasons sampled. However, PAC-contaminated sediments in the Athabasca Delta and Lake Athabasca (35, 36) are consistent with long-range atmospheric and fluvial transport of particulate PAC. Our sampling did not include the intervening spring snowmelt, which would release a pulse of PAC up to 50 km from oil sands upgrading facilities into nearby tributaries and the Athabasca. Although this should increase PAC concentrations in tributaries, it might not be detected in the main stem because of high-volume dilution by uncontaminated snowmelt from areas upstream of oil sands mining. PAC may also be removed from the water column by microbial degradation and adsorption to organic matter that settles out of the water column (37). These dilution and removal processes likely account for declining con-

centrations of dissolved PAC as the Athabasca flows to Lake Athabasca and Fort Chipewyan.

## Conclusions

Due to substantial loadings of airborne PAC, the oil sands industry is a far greater source of regional PAC contamination than previously realized. Despite previous recommendations (17), there is no apparent detailed monitoring of PAC fluxes via wet and dry deposition in the winter or summer, when similar or greater contributions are likely. Monitoring of air, the snowpack, spring snowmelt, and summer rain and vegetation is essential to identify and control sources of PAC and their potential environmental and human health impacts. A second important source of PAC is landscape disturbance. Surprisingly, impacts are related primarily to recent disturbance (<2 y), which suggests that revegetation or erosion controls mitigate long-term loadings.

Controls on waterborne PAC are critical because concentrations at tributary mouths and at one site on the Athabasca are already within the range toxic to fish embryos. However, the impacts on the Athabasca ecosystem of mining wastewater, snowmelt, or contaminated groundwater remain enigmatic due to high seasonal variability of flow and dilution capacity.

Our study confirms the serious defects of the RAMP (12). More than 10 years of inconsistent sampling design, inadequate statistical power, and monitoring-insensitive responses have missed major sources of PAC to the Athabasca watershed. Most importantly, RAMP claims that PAC concentrations are within baseline conditions and of natural origin have fostered the perception that high-intensity mining and processing have no serious environmental impacts. The existing RAMP must be redesigned with more scientific and technical oversight to better detect and track PAC discharges and effects. Oversight by an independent board of experts would make better use of monitoring resources and ensure that data are available for independent scrutiny and analyses. The scale and intensity of oil sands development and the complexity of PAC transport and fate in the Athabasca watershed demand the highest quality of scientific effort.

## Methods

**Study Design.** Seventeen sites were chosen on the main stem Athabasca, the Delta, and Lake Athabasca, from south of Fort McMurray (upstream of development) to Fort Chipewyan (downstream of development) (Fig. 1). The Athabasca River is exposed to the McMF 50 km upstream of Fort McMurray and is present within its banks to Eymundson Creek (38).

Tributaries draining from the east (Steeptank, Muskeg, and Firebag Rivers) and west (Beaver, Ells, and Tar Rivers), including reference rivers (Firebag and Ells Rivers) and those impacted by oil sands development (Steeptank, Muskeg, Beaver, and Tar Rivers), and the Horse River stream mouth, were sampled in winter and summer (Fig. 1). Nine other stream mouth sites (AR17down, AR17up, Clarke, Poplar, McLean, Fort, and Eymundson Creeks and MacKay and Calumet Rivers) were sampled only in summer.

GIS analyses were performed by using ArcGIS 9 ArcMap version 9.2 (39), to delineate catchments, extract disturbance and geologic data, and calculate distances between AR6 and other snow sampling sites (see *Analytical and Statistical Method Details* in *SI Text* for details).

**Field Sampling.** In March, snow was collected from 12 sites on the Athabasca River, Athabasca Delta, and Lake Athabasca and from 19 tributary sites. Samples were collected close to the middle of the river, and replicated at MU1. At each site, the depth and weight of five snow cores were recorded to calculate snow density, snow water equivalents, and PAC areal deposition rates. For PAC, an integrated sample of the snowpack was melted and vigorously stirred, and a subsample (775–4,000 mL) was filtered through a  $0.45\text{-}\mu\text{m}$  muffled Whatman GF/F filter. The filter was frozen, and the filtrate was decanted, spiked with a suite of perdeuterated hydrocarbon surrogate standards in 0.5 mL of acetone, shaken, extracted twice with 100 mL of DCM, and stored at  $-20^\circ\text{C}$ .

PMDs were deployed in the Athabasca and its tributaries for  $\approx 30$  days to passively monitor dissolved PAH in summer and winter and provide a time-integrated measure of dissolved PAH (14). PMDs were cleaned by sonication successively with dichloromethane (DCM), put in DCM-cleaned metal holders,

wrapped in DCM-rinsed aluminum foil and stored in heat-sealed Ziploc bags at  $-20^{\circ}\text{C}$ . At 12 locations on the Athabasca River and the Horse River mouth, two PMDs were deployed per site, near the river bottom and in surface water. One PMD was deployed near the river bottom at the five other Athabasca sites, and all tributary sites. PMDs were retrieved by using global positioning system coordinates and a metal detector (winter). Replicate PMDs were deployed within 500 m at sites where PAC were assumed to be high (MU3) or low (EL1). One trip blank and five field blanks, handled like deployed PMDs, were included with winter and summer samples.

Oil sands samples were collected in summer from the Syncrude lease area, the east and west bank of the Athabasca River north of Bitumount, and underwater at the mouth of the Steepbank River. All samples were frozen.

**Laboratory.** PAC were measured at the University of Alberta Biogeochemical Analytical Laboratory by using an Agilent 6890N gas chromatograph coupled to a 5975 inert XL mass selective detector and 7683B injector (modified refs. 14 and 40). Detailed methods and instrument and method detection limits are

presented in *Analytical and Statistical Method Details* in *SI Text* and *Table S1*, respectively.

**Analytical and Statistical Methods.** Details of QA/QC, contamination by diesel fuel, PAC source identification, estimation of aqueous PAC concentrations and PAC deposition calculations are provided in *Analytical and Statistical Method Details* in *SI Text*.

**ACKNOWLEDGMENTS.** Dr. K. Timoney, R. Grandjambe, R. Radmanovich, R. Ladouceur, S. Laurent, T. Boag, P. Jordan, K. Vladicka-Davies, M. Davies, and H. La provided assistance with planning and field and/or laboratory work. Parks Canada and the Government of Alberta granted research permits. The Athabasca Chipewyan, Mikisew Cree, and Fort MacKay First Nations and Métis provided traditional ecological knowledge and access to traditional territories. Matt Hanneman and Peter Lee (Global Forest Watch) supplied GIS data and defined catchments. Charlene Nielsen completed GIS analyses and created the study area map. Reviewer comments improved the manuscript. This work was funded by the Tides Foundation and the Walter and Duncan Gordon Foundation.

- Government of Alberta (2008) *Environmental Management of Alberta's Oil Sands*. (Oil Sands Management Division, Government of Alberta, AB, Canada).
- ERCB (Energy Resources Conservation Board) (2005) *Alberta's Reserves 2004 and Supply/Demand Outlook 2005–2014*. ST98-2005, Power Point file of graphs and data from 2005 report. Available from [www.ercb.ca/portal/server.pt/gateway/PTARGS.0.0.308.265.0.43/http%3B/ercbContent/publishedcontent/publish/ercb.home/publications.catalogue/publications.available/serial/publications/st98.aspx](http://www.ercb.ca/portal/server.pt/gateway/PTARGS.0.0.308.265.0.43/http%3B/ercbContent/publishedcontent/publish/ercb.home/publications.catalogue/publications.available/serial/publications/st98.aspx).
- ERCB (Energy Resources Conservation Board) (2009) *Alberta's Energy Reserves 2008 and Supply/Demand Outlook 2009–2018*. ST98-2009, Power Point file of graphs and data from 2009 report. Available from: [www.ercb.ca/portal/server.pt/gateway/PTARGS.0.0.308.265.0.43/http%3B/ercbContent/publishedcontent/publish/ercb.home/publications.catalogue/publications.available/serial/publications/st98.aspx](http://www.ercb.ca/portal/server.pt/gateway/PTARGS.0.0.308.265.0.43/http%3B/ercbContent/publishedcontent/publish/ercb.home/publications.catalogue/publications.available/serial/publications/st98.aspx).
- Price M (2008) *11 Million Litres a Day the Tar Sands' Leaking Legacy*. (Environmental Defense, Toronto, ON, Canada).
- CAPP (Canadian Association of Petroleum Producers) (2009) *Crude Oil Forecast, Markets and Pipeline Expansions* (Canadian Association of Petroleum Producers).
- Chen Y (2009) *Cancer Incidence in Fort Chipewyan, Alberta 1995–2006* (Alberta Cancer Board, Division of Population Health and Information Surveillance, Alberta Health Services).
- RAMP (Regional Aquatics Monitoring Program), WBEA (Wood Buffalo Environmental Association), CEMA (Cumulative Environmental Management Association) (2008) *Joint Community Update 2008 Reporting Our Environmental Activities to the Community* (RAMP, WBEA and CEMA, Fort McMurray, AB, Canada).
- HEAP (The Alberta Oil Sands Community Exposure and Health Effects Assessment Program) (2000) *Summary Report. Health Surveillance* (Alberta Health and Wellness, Government of Alberta, Edmonton, AB, Canada), [www.health.alberta.ca/newsroom/pub-environmental-health.html](http://www.health.alberta.ca/newsroom/pub-environmental-health.html).
- HEMP (Wood Buffalo Environmental Association Human Exposure Monitoring Program) (2007) *Methods Report and 2005 Monitoring Year Results* (Wood Buffalo Environmental Monitoring Association, Fort McMurray, AB, Canada), [www.health.alberta.ca/newsroom/pub-environmental-health.html](http://www.health.alberta.ca/newsroom/pub-environmental-health.html).
- HEMP (Wood Buffalo Environmental Association Human Exposure Monitoring Program) (2007) *2006 Monitoring Year Results* (Wood Buffalo Environmental Monitoring Association, Fort McMurray, AB, Canada), [www.health.alberta.ca/newsroom/pub-environmental-health.html](http://www.health.alberta.ca/newsroom/pub-environmental-health.html).
- RAMP (Regional Aquatics Monitoring Program) (2009A) *2008 Final Technical Report* (Hatfield Consultants, Kilgour and Associates Ltd., Klohn Crippen Berger Ltd., and Western Resource Solutions).
- Ayles GB, Dubé M, Rosenberg D (2004) *Oil Sands Regional Aquatic Monitoring Program (RAMP) Scientific Peer Review of the Five Year Report (1997–2001)* (Fisheries and Oceans Canada, Winnipeg, Manitoba, Canada).
- Strausz OP, Lown EM (2003) *The Chemistry of Alberta Oil Sands, Bitumens and Heavy Oils* (Alberta Energy Research Institute, Calgary, AB, Canada).
- Carls MG, Holland LG, Short JW, Heintz RA, Rice SD (2004) Monitoring polynuclear aromatic hydrocarbons in aqueous environmental with passive low-density polyethylene membrane devices. *Environ Toxicol Chem* 23:1416–1424.
- Wakeham SC, Schaffner C, Giger W (1980) Polycyclic aromatic hydrocarbons in recent lake sediments I. Compounds having anthropogenic origins. *Geochim Cosmochim Acta* 44:415–429.
- Barrie LA, Kovalick J (1980) *A Wintertime Investigation of the Deposition of Pollutants Around an Isolated Power Plant in Northern Alberta*. Prepared for the Alberta Oil Sands Environ Res Program by Atmos Environ Service. AOSERP Report 90 (Alberta Oil Sands Environmental Research Program).
- Murray WA (1981) The 1981 snowpack survey in the AOSERP study area. Prepared for the Alberta Oil Sands Environ Res Program by Promet Environmental Group. AOSERP Report 125 (Alberta Oil Sands Environmental Research Program).
- National Pollutant Release Inventory (NPRI) (2009) [www.ec.gc.ca/inrp-npri/](http://www.ec.gc.ca/inrp-npri/).
- Allan R, Jackson TA (1978) Heavy metals in bottom sediments of the mainstem Athabasca River system in the AOSERP study area. Prepared for the Alberta Oil Sands Environ Res Program by Inland Waters Directorate, Fisheries and Environment Canada. AOSERP Report 34 (Alberta Oil Sands Environmental Research Program).
- Yunker MB, Backus SM, Pannatier EG, Jeffries DS, Macdonald RW (2002) Sources and significance of alkane and PAH hydrocarbons in Canadian arctic rivers. *Estuar Coast Shelf Sci* 55:1–31.
- Carls MG, Rice SD, Hose JE (1999) Sensitivity of fish embryos to weathered crude oil: Part I. low-level exposure during incubation causes malformations, genetic damage, and mortality in larval pacific herring (*Clupea pallasii*). *Environ Toxicol Chem* 18:481–493.
- Hodson PV, et al. (2007) Alkyl PAH in crude oil cause chronic toxicity to early life stages of fish. *Proceedings of the 28th Arctic and Marice Oilspill Program (AMOP)* (Technical Seminar, Environmental Science and Technology Division, Environment Canada, Ottawa, ON, Canada), pp 291–300.
- Incardona JP, Collier TK, Scholz NL (2004) Defects in cardiac function precede morphological abnormalities in fish embryos exposed to polycyclic aromatic hydrocarbons. *Toxicol Appl Pharmacol* 196:191–205.
- Colavecchia MV, Backus SM, Hodson PV, Parrott JL (2004) Toxicity of oil sands to early life stages of fathead minnows (*Pimephales promelas*) *Environ Toxicol Chem* 23:1709–1718.
- Colavecchia MV, Hodson PV, Parrott JL (2006) CYP1A induction and blue sac disease in early life stages of white sucker (*Catostomus commersoni*) exposed to oil sands. *J Toxicol Environ Health A* 69:967–994.
- Tetreault GR, McMaster ME, Dixon DG, Parrott JL (2003) Using reproductive endpoints in small forage fish species to evaluate the effects of Athabasca oil sands activities. *Environ Toxicol Chem* 22:2775–2782.
- Timoney K (2007) *A Study of the Water and Sediment Quality as Related to Public Health Issues, Fort Chipewyan, Alberta*. (Treeline Ecological Research, Edmonton, AB, Canada).
- Sharma M, McBean EA (2001) PAH deposition to snow surface. *Environ Sci Pollut Res* 8:11–18.
- Meyer T, Wania F (2008) Organic contaminant amplification during snowmelt. *Water Res* 42:1847–1865.
- RAMP (Regional Aquatics Monitoring Program) (2009B) *Appendices. 2008 Technical Report* (Hatfield Consultants, Kilgour and Associates Ltd., Klohn Crippen Berger Ltd., and Western Resource Solutions).
- Scott WB, Crossman EJ (1998) *Freshwater Fishes of Canada* (Galt House Oakville, ON, Canada).
- Newstedt JL, Giesy JP (1987) Predictive models for photoinduced acute toxicity of polycyclic aromatic hydrocarbons to *Daphnia magna*, Strauss (Cladocera, Crustacea). *Environ Toxicol Chem* 6:445–461.
- Vehniainen ER, Hakkinen J, Oikari A (2003) Photoinduced lethal and sublethal toxicity of retene, a polycyclic aromatic hydrocarbon derived from resin acid, to coregonid larvae. *Environ Toxicol Chem* 22:2995–3000.
- Xie F, et al. (2006) Assessment of the toxicity of mixtures of copper, 9,10-phenanthrenequinone, and phenanthrene to *Daphnia magna*: Evidence for a reactive oxygen mechanism. *Environ Toxicol Chem* 25:165–174.
- Evans M (2003) Assessment of natural and anthropogenic impacts of oil sands contaminants within the northern river basins, Long-range Transport of Hydrocarbons to the Northern Deltas and Lakes: Pathways and Fate. *Assessment of Natural and Anthropogenic Impacts of Oil Sands Contaminants Within the Northern River Basins - Final Summary Report - Task 5: Hydrocarbons/oil sands and heavy oil research and development*, Brua RB, Cash KL Culp JM. Submitted to the Panel on Energy Research and Development (Environment Canada, Saskatoon, Saskatchewan, Canada).
- Akre CJ, Headley JV, Conly FM, Peru KM, Dickson LC (2004) Spatial patterns of natural polycyclic aromatic hydrocarbons in sediment in the lower Athabasca River. *J Environ Health A* 39:1163–1176.
- French-McCay DF (2004) Oil spill impact modeling: Development and validation. *Environ Toxicol Chem* 23:2441–2456.
- Conly FM, Crosley RW, Headley JV (2002) Characterizing sediment sources and natural hydrocarbon inputs in the lower Athabasca River, Canada. *Environ Eng Sci* 1:187–199.
- ESRI (Environmental Systems Research Institute) (2006) *ArcGIS: Release 9.2* (Environmental Systems Research Institute, 1999–2006, Redlands, CA), [www.esri.com](http://www.esri.com).
- Short JW, Jackson TJ, Larsen ML, Wade TL (1996) Analytical methods used for the analysis of hydrocarbons in crude oil, tissues, sediments, and seawater collected for the natural resources damage assessment for the Exxon Valdez oil spill. *Am Fish Soc Symp* 18:140–148.